
Structural comparison of yeast ribosomal protein genes

Robert J. Leer, Mary M.C. van Raamsdonk-Duin, Mark J.M. Hagendoorn, Willem H. Mager and Rudi J. Planta

Biochemisch Laboratorium, Vrije Universiteit, de Boelelaan 1083, 1081 HV Amsterdam, The Netherlands

Received 2 July 1984; Revised and Accepted 7 August 1984

ABSTRACT

The primary structure of the genes encoding the yeast ribosomal proteins L17a and L25 was determined, as well as the positions of the 5'- and 3'-termini of the corresponding mRNAs. Comparison of the gene sequences to those obtained for various other yeast ribosomal protein genes revealed several similarities. In all split genes the intron is located near the 5'-side of the amino acid coding region. Among the introns a clear pattern of sequence conservation can be observed. In particular the intron-exon boundaries and a region close to the 3'-splice site show sequence homology. Conserved sequences were also found in the leader and trailer regions of the ribosomal protein mRNAs. The 5'-flanking regions of the yeast ribosomal protein genes appeared to contain sequence elements that many but not all ribosomal protein genes have in common, and therefore may be implicated in the coordinate expression of these genes. The amino acid coding sequences of the ribosomal protein genes show a biased codon usage. Like most yeast ribosomal protein molecules, L17a and L25 are particularly basic at their N-terminus.

INTRODUCTION

The biogenesis of ribosomes is a highly complex process requiring the coordinate and balanced synthesis of a large number of constituents. In *E. coli* the details of the regulation mechanisms underlying this coordinate synthesis are beginning to emerge. As far as synthesis of ribosomal proteins is concerned, it has become clear that the organization of ribosomal protein genes into a limited number of transcriptional and translational units is an important feature of the regulation mechanism (see Ref. 1 for a recent review). Regulation of ribosomal protein synthesis in eukaryotic cells so far is by and large terra incognita. The best-studied organism in this field is yeast, where about 30 different ribosomal protein genes have so far been cloned by ourselves as well as others (2-8). In our group we have isolated the genes for proteins S10, S16A, S24, S31, S33, rp28, L16, L17a, L25, L34, L36 and L46 from a colony bank of HindIII-generated yeast DNA fragments cloned in pBR322 (8). Genomic blotting experiments showed that almost all of these genes are duplicated, the gene for L25 being the only exception (9). Moreover, both our

own studies and experiments by others have revealed that yeast ribosomal protein genes in general are not clustered (3,8,10). The only exceptions to this rule so far are the genes encoding proteins S16A and rp28, which are separated by only about 600 bp (Molenaar, C.M.T. and Pearson, N.J., manuscript in preparation) and those encoding S24 and L46 which are about 700 bp apart (R.J. Leer, manuscript in preparation).

Ribosomal protein genes in yeast are somewhat of a class apart since most have been found to contain an intron (3,8,9,11). So far only four ribosomal protein genes [L3 (6), L16 (24), S33 (12) and S24 (R.J. Leer, manuscript in preparation)] do not conform to this rule. In other yeast genes the presence of an intron is the exception rather than the rule. Out of a large number of genes studied as yet only two [the actine gene (13,14) and the MAT1 α -gene (15)] are split.

The split nature of the ribosomal protein genes explains the coordinate depression of ribosomal protein synthesis in splicing-defective ts mutants of yeast upon raising the temperature (16,17). We (9) and others (3,11) have shown that under those conditions pre-mRNAs for several ribosomal protein species accumulate that are 300-500 nucleotides longer than their mature counterparts. The ribosomal proteins known to escape this coordinate regulation are precisely the ones mentioned above as containing no intron in their gene. Clearly processing of mRNA is an important regulatory mechanism under restrictive conditions in these ts-mutants.

In wild-type yeast cells coordinate control of ribosomal protein synthesis may take place at both the transcriptional and translational level. For instance, after a temperature shock a sudden coordinate decrease in ribosomal protein gene transcription occurs (11,18). On the other hand the outcome of gene dosage experiments using yeast ribosomal protein genes cloned in multi-copy vectors, indicate that control is also exerted at the translational level (19; J.R. Warner, personal communication).

In order to classify further the molecular mechanisms regulating the coordinate and balanced synthesis of ribosomal proteins in yeast we are carrying out a detailed structural analysis of a number of individual yeast ribosomal protein genes. The aim of these studies is the identification of common sequences that might act as signals in regulatory events. Deletion mapping and site-directed mutagenesis can then be used to ascertain the function of these sequences. In view of the evidence for both transcriptional and translational regulation we have focussed our attention on the untranslated mRNA sequences as well as the DNA sequences lying upstream of the tran-

scription initiation site of the ribosomal protein genes. In this paper we present such an analysis of the genes for the large subunit proteins L17a and L25 and compare the results with previously published data on several yeast ribosomal protein genes.

MATERIALS AND METHODS

DNA preparation and sequence analysis

Plasmids pBMCY111 and pBMCY138 containing the genes coding for L17a and L25 respectively, were purified as described previously (12). Restriction enzyme digestions were performed as recommended by the suppliers (New England Laboratories; Boehringer Mannheim). The cutting sites for the enzymes TaqI, MspI and HaeIII were mapped using the partial digestion procedure of Smith and Birnstiel (20). DNA fragments were subcloned in M13 mp7, mp8 or mp9 and their sequence was determined by the dideoxy chain termination method (21,22). Sequence analysis according to the chemical procedure was performed as described in (23).

Primer extended sequence analysis of mRNA

Details of primer preparation and RNA sequencing using reverse transcriptase are described elsewhere (24). The primer for the L17a gene was labelled by repair synthesis of the HaeIII-BglII fragment in M13 mp8 (24) (see Fig. 1). After digestion with RsaI a fragment of 107 nucleotides was isolated (24 nucleotides of which originate from the M13 cloning region) (24). The primer for the L25 gene was a 75 nucleotides long DNA fragment obtained by SacI + RsaI digestion of cDNA synthesized using the subcloned HpaI-TaqI fragment in M13 mp9 as a template (24) (see Fig. 2).

S1 nuclease mapping of the 3'-ends of mRNAs

Preparation of the probes and S1 nuclease analysis were carried out as described in detail elsewhere (24). The probe for the transcript of the L17a gene was a 316 nucleotides long DNA fragment obtained by HindIII digestion of cDNA synthesized using the subcloned Taq-Taq fragment in M13 mp9 as a template (24) (see Fig. 1; 47 nucleotides originate from M13).

For mapping the 3'-end of the L25 gene transcript pBMCY138 DNA was digested with HpaI. After limited treatment with exonuclease III and repairing the 3'-end with Klenow polymerase in the presence of α -³²P-dATP, the 264 nucleotides long probe was isolated as described elsewhere (24).

RESULTS AND DISCUSSION

From the colony bank of HindIII-generated yeast DNA fragments in pBR322 that has been described in a previous paper (4), a number of ribosomal protein




Fig. 1. Map of the insert of pBMCY111 and structural analysis of the L17a gene.

The position of some restriction enzyme sites as well as the location of the R-loop indicated in A were published previously (8). H = HindIII, S = Sau3A, M = MspI, T = TaqI, R = RsaI (only RsaI sites are indicated that are used for constructing the primers), Ha = HaeIII, B = BglIII, K = KpnI.

In B the sequence strategy is shown. The arrows give the extent of nucleotide analysis according to the chain termination method; the dashed arrow indicates a sequence determined using the chemical procedure. The 269 nucleotides long probe used for the 3'-mapping experiment and the 73 nucleotides long primer used for the primer extension experiment are indicated. The structure of the L17a transcript is given in C.

genes has been isolated (8). Recombinant DNA pBMCY111 carries a gene coding for ribosomal protein L17a (8; R.J. Leer, unpublished results) whereas clone pBMCY138 contains the gene encoding the large subunit protein L25 (8). Southern analysis using genomic digests of yeast DNA revealed that L25 is encoded by a simple locus (data not shown; see Ref. 9) whereas the gene encoding L17a is duplicated on the yeast genome (9).

In Figures 1 and 2 the physical maps of the respective cloned DNA fragments are depicted. The genes were located in the maps by electron microscopic R-loop analysis (8). The strategy followed in sequencing the respective genes is also outlined.

The results of the nucleotide sequence analysis are shown in Figures 3 and 4. The open reading frame of the L17a gene encompasses 411 nucleotides, coding for a protein with a molecular mass of 14,321 D and a net charge of +13. The gene for L25 consists of 414 nucleotides encoding a protein with a molecular mass of 15,225 D and a net charge of +19. The identity of the protein encoded by the L25 gene is confirmed by the amino acid composition deduced from the nucleotide sequence. This composition is in good agreement with corrected data obtained from studies on protein YP42' = YL25 (25; E. Otaka, personal communication) which is identical to L25 (26). No such data so far exist for protein L17a, which is identical to YL32 (26). It is noteworthy that both L25 and L17a are particularly basic at their N-terminus. Neither protein contains acidic residues within the first 55 amino acids. A similar




Fig. 2. Map of the insert of pBMCY138 and structural analysis of L25 gene. The position of some restriction enzyme sites as well as the location of the R-loop indicated in A were published previously (8). For restriction enzyme sites the same symbols are used as in Fig. 1. In addition Sc = SacI, Hp = HpaI, E = EcoRI. In B the sequence strategy is shown. The 264 nucleotides long probe for the 3'-mapping experiment as well as the 75 nucleotides long primer for cDNA synthesis are indicated. The sequence spanning over the Sau3A site at the 5' end of the gene was confirmed by primer extension analysis. In C the structure of the L25 transcript is given.

skewed distribution of charged amino acids is observed in several other, though not all yeast ribosomal proteins. Furthermore L25, which probably is one of the primary assembling ribosomal proteins (27) is characterized by the presence of 5 Lys-Lys doublet sequences, 4 of which are part of a Lys-Lys-Ala tripeptide.

The amino acid coding sequence of the genes for L17a and L25 reveals a biased codon usage, in agreement with findings for other yeast ribosomal protein genes (see Table I). The pattern of preferred codons in ribosomal protein genes is quite similar to the one observed for highly expressed yeast genes (28) and may reflect an efficient translation of the moderately abundant ribosomal protein mRNAs.

As can be concluded from the sequence data presented in Figures 3 and 4 the open reading frames of both the L17a and the L25 gene are interrupted by an intron of 513 and 415 nucleotides respectively. In this respect the genes for L17a and L25 are similar to most other ribosomal protein genes in yeast. A remarkable feature of all split yeast ribosomal protein genes studied so far is the location of the intervening sequence close to the 5'-end of the gene (cf. Table II). Recent findings (9,35) indicate that in some ribosomal protein genes the intron is present even upstream of the ATG codon. Since a split character is rarely found for non-ribosomal protein genes on the yeast nuclear genome it is tempting to suggest a regulatory function for these introns. Whether the interruption of many yeast ribosomal protein genes at their very 5'-end is implicated in transcriptional or post-transcriptional control, however, is still open to speculation.

```

-420          -400          -380          -360          -340
AAGCTTGAATACCAATCACTATTACCATGTTAAATGATCCCCAGATAAAAAAGACAAGACAGCGGTATATCAGACTACTTTAACATCTGTACACCCCTCA

-320          -300          -280          -260          -240
ATTGCTTTGAAAAAACCCATGCAGCTATTATATTTTTTCAACGGACCAATTTTTTTTATCGCTTATCCGGCTCACAGAAACGAGCCCTACAACCGTCAATCAG

-220          -200          -180          -160          -140
ACTTGCATGGAATGTTGAAAAAGAGGTAATAAAGAAAAATATCAAAACCAACACACTCGGGCTGAAAAATTAATATGCGAGAAACGAACTGTAGCCAT

-120          -100          -80          -60          -40
AATGTACCTGTTCCCGCTAATAGTAATACCATCGTATTGCTAATGTTAGTAGCTAAAAATCTCTTACTTCTTTATTTTGATTCTTGGGCTTACAAGCA

ATACAAAACCAACACACACTATATATAACTAATA ATG TCC GGT AAC GGT GCT CAA GGT ACT AAG TTT AGA ATC TCA GTATCTTAAA
      ↓   ↓
      met ser gly asn gly ala gln gly thr lys phe arg ile ser

60          80          100          120          140
ATTTTTATTTTCCACAATGCAATTTGGTTAAATTTGATCATAAAGTAAAGTTCCAAGATTTTCATTTGCTGGGTACAACAGAATTAACAGAGGTTTAAAA

160          180          200          220          240
GACGCTAAGACGCTAAACACTAAGTGTGAAGCCGTATCATATACTTATCGATGTTATTTCCATAAGAATGCTAGATGTCCATCACTGTTACCGCCCCG

260          280          300          320          340
CCAAAGATGCAATTACATACATAAACTGAGTTTTTAAATGGTTAAGCGGACTGTATTGTCAGATGAAGTGATACCTGTCAATATTCATGGACTCAAAAAAT

360          380          400          420          440
TTCTGTTTTCAAAATCATAACCGCAAGCATATCTCTAACTCCCTCTTTTGCCCATACTAATGGATAACCATCGAAATGTTATTTTACAAAAATCACACT

460          480          500          520          540
TAACTCTATTTTCAAGATTAACACATCTGCTAACTTTTCGCCATGCTTTTACTAACAAAATTAATCTCACTCCCGATATTGAGATCTTTTTTAACT

560          580          600          620
AG TTA GGT CTA CCA GTC GGT GCC ATC ATG AAC TGT GCT GAC AAC AGT GGT GCC AGA AAC TTG TAC ATT ATC GCC
  leu gly leu pro val gly ala ile met asn cys ala asp asn ser gly ala arg asn leu tyr ile ile ala

640          660          680          700
GTC AAA GGC TCT GGT TCC AGA TTG AAC AGA TTG CCA GCC GCC TCT CTA GGT GAT ATG GTT ATG GCC ACC GTT AAG
val lys gly ser gly ser arg leu asn arg leu pro ala ala ser leu gly asp met val met ala thr val lys

720          740          760
AAG GGT AAG CGA GAA TTG AGA AAG AAG GTT ATC CCA GCT ATT GTT GTC CGT CAA GGT AAG TCT TGG AGA AGA AGA
lys gly lys pro glu leu arg lys lys val met pro ala ile val val arg gln ala lys ser trp arg arg arg

780          800          820          840
GAC GGT GTC TTT TTG TAC TTC GAA GAC AAT GCT GGT GTC ATC GCT AAT CCT AAG GGT GAA ATG AAG GGT TCC GCC
asp gly val phe leu tyr phe glu asp asn ala gly val ile ala asn pro lys gly glu met lys gly ser ala

860          880          900          920
ATC ACT GGT CCA GTC GGT AAG GAA TGT GCC GAT TTA TGG CCA AGA GTT GGA TCT AAC TCC GGT GTT GTT GTG TAA
ile thr gly pro val gly lys glu cys ala asp leu trp pro arg val ala ser asn ser gly val val val end

940          960          980          1000          1020
AAAGGAAATATTTTGACTCATACTCTTAGATCTTCAATTATATATATTTTCTTTATAAATAACAATCTCTTAAACAGATTAAGAAAACTTAAATATTC
      ↓
ATTGCATTTTATGATGAAAAATATTACTATTATCGTTTCATCGAGAGAATCACATGATATTATGAGCGTGATAGGAGGCTTATATATTTTTGGGGTAGGT

1040          1060          1080          1100          1120
CAAAAAAGCAAAACAGACACGTTAAGAAGCTAAGAATAACACAGATGATAGAATGTTGTGCTAGTGAGAAAACAGTATGCCACCGCCGTACAATATGCGTATT

1140          1160          1180          1200          1220
TTGACATTTTCAAGGTAATAGTTCCAAGTTAATAGAATAATCAAGACTAATTGACGTCTCAAAGCCTCGCAAAAAAAGCAGGAAATTCGAAAAAATAA

1240          1260          1280          1300          1320
GCCTTCAAAGAAAACGGAAATCACTGTAACATAGACGATGGAAGATCATCGGATTGTAGAACTTTAAACTTTCTATCAACAAAAATCAAGAGAGA

1340          1360          1380          1400          1420
AACAAATGCTTTAGATGACGTAACAACAATTTTAAAGAAGATCCGAAAAGGATACCAACCAAGGCCATATCTACAACCGCAGAGCTT

```

Fig. 3. DNA sequence of the gene for yeast ribosomal protein L17a and amino acid sequence of the L17a protein. The nucleotides are numbered with italics starting from the first nucleotide of the initiation codon. The positions of the 5'- and 3'-ends of the L17a mRNA are indicated by arrows.




Fig. 4. DNA sequence of the gene for yeast ribosomal protein L25 and amino acid sequence of the L25 protein.

The nucleotides are numbered with italics starting from the first nucleotide of the initiation codon. The positions of the 5'- and 3'-ends of the L25 mRNA are indicated by arrows.

Table I. Codon usage for the genes coding for yeast ribosomal proteins L17a and L25 as compared with other ribosomal protein genes (RP) and several highly expressed yeast genes (Y).

	L17a		L25		RP ^{a)}		Y ^{b)}		L17a		L25		RP ^{a)}		Y ^{b)}							
	2	1	2	1	3	4	4	3	46	79	73	73	38	73	5	2						
Phe UUC	2	-	2	5	Ser	UCU	4	3	46	79	Tyr	UAU	-	1	5	2	Cys	UGU	2	-	6	20
Phe UUC	1	2	52	63	Ser	UCC	4	2	29	73	Tyr	UAC	2	7	38	73	Cys	UGC	-	-	1	2
Leu UUA	2	1	13	14	Ser	UCA	1	1	3	1												
Leu UUG	5	8	94	148	Ser	UCG	-	-	-	1							Trp	UGG	2	-	11	26
Leu CUU	-	-	1	-	Pro	CCU	1	-	5	6	His	CAU	-	-	12	3	Arg	CGU	1	-	25	2
Leu CUC	-	-	-	1	Pro	CCC	-	-	1	1	His	CAC	-	1	30	56	Arg	CGC	-	-	-	-
Leu CUA	2	1	8	6	Pro	CCA	6	6	52	71	Gln	CAA	2	3	50	44	Arg	CGA	-	-	-	-
Leu CUG	-	-	-	-	Pro	CCG	-	-	-	-	Gln	CAG	-	-	1	-	Arg	CGC	-	-	-	-
Ile AUU	2	5	39	58	Thr	ACU	2	5	46	49	Asn	AAU	2	1	3	3	Ser	AGU	1	-	1	1
Ile AUC	5	3	42	66	Thr	ACC	1	4	38	66	Asn	AAC	6	6	46	86	Ser	AGC	-	-	4	1
Ile AUA	-	-	-	1	Thr	ACA	-	-	2	-	Lys	AAA	1	4	26	21	Arg	AGA	9	7	135	63
Met AUG	6	3	28	39	Thr	ACG	-	-	-	-	Lys	AAG	10	17	141	151	Arg	AGG	-	-	1	1
Val GUU	7	7	73	112	Ala	GCU	6	17	90	182	Asp	GAU	2	2	33	37	Gly	GGU	13	5	120	164
Val GUC	6	5	66	93	Ala	GCC	8	2	28	62	Asp	GAC	3	3	40	100	Gly	GGC	1	-	4	11
Val GUA	-	-	-	3	Ala	GCA	1	-	4	3	Glu	GAA	4	4	89	67	Gly	GGA	-	-	2	1
Val GUG	1	-	2	1	Ala	GCG	-	-	2	4	Glu	GAG	-	1	1	3	Gly	GGG	-	-	1	1

a) Ribosomal protein genes encoding S10 (29), S33 (12), rp51 (30), rp28 and S16A (Molenaar, C.M.T. and Pearson, N.J., manuscript in preparation), L16 (24), L34 (31), L3 (6) and L29 (32).
 b) From Ref. 33.

Table II. General structure of yeast ribosomal protein genes.

		5' exon			intron (a)		3' exon		
		start	no. of codons		no. of nucleotides		no. of codons	stop	
S10	(b)	ATG	-	1	←	352 (394)	→	234	- TAA
L34	(c)	ATG	-	18	←	349 (421)	→	94	- TAA
S16A	(d)	ATG	-	5	←	390 (551)	→	138	- TAA
rp28	(d)	ATG	-	36	←	429 (447)	→	149	- TAA
L46	(e)	ATG	-	1	←	383	→	49	- TAA
rp51	(f)	ATG	-	-	←	325 (398)	→	135	- TAA
L29	(g)	ATG	-	15	←	510	→	133	- TAA
L17a		ATG	-	13	←	513	→	123	- TAA
L25		ATG	-	3	←	415	→	134	- TAA

- (a) data for the second gene copy are within brackets;
- (b) (29) and R.J. Leer, unpublished results;
- (c) (31);
- (d) Molenaar, C.M.T. and Pearson, N.J., manuscript in preparation;
- (e) R.J. Leer, unpublished results;
- (f) (30,34);
- (g) (32)

The introns occurring in yeast ribosomal protein genes show several sequence similarities as outlined in Table III. The structures of the 5'- and 3'-splice sites without exception are in perfect agreement with the consensus boundary

Table III. Intron sequences within yeast ribosomal protein genes (a).

S10-1	↓	GTATGT	-	316 n.	-	TTTACTAACA	-	10 n.	-	TTTATAACAG	↓
S10-2		GTATGA	-	343 n.	-	TATACTAACA	-	25 n.	-	TTTTAAACAG	
L34-1		GTATGT	-	378 n.	-	GTTACTAACA	-	17 n.	-	TTTTTAATAG	
L34-2		GTATGT	-	305 n.	-	TTTACTAACA	-	18 n.	-	ATTAATATAG	
S16A-1		GTACGT	-	341 n.	-	TTTACTAACA	-	23 n.	-	TTTTCTACAG	
S16A-2		GTACGT	-	503 n.	-	TTTACTAACA	-	22 n.	-	TTTCAATTAG	
rp28-1		GTATGT	-	409 n.	-	GTTACTAACA	-	12 n.	-	TTTTTTTTAG	
rp28-2		GTATGT	-	375 n.	-	TTTACTAACA	-	28 n.	-	TTAATCACAG	
L46		GTATGT	-	314 n.	-	GTTACTAACA	-	43 n.	-	TTTTAATTAG	
rp51-1		GTATGT	-	326 n.	-	TATACTAACA	-	46 n.	-	ATTTAATAG	
rp51-2		GTACGT	-	282 n.	-	TTTACTAACT	-	23 n.	-	ATTTTTATAG	
L29		GTATGT	-	453 n.	-	TTTACTAACG	-	31 n.	-	TTTTGTACAG	
L17a		GTATGT	-	455 n.	-	TTTACTAACA	-	32 n.	-	TTTTAACTAG	
L25		GTATGT	-	360 n.	-	TTTACTAACA	-	29 n.	-	TTTTGTATAG	
consensus	↓	GTA ^T _C Gt	-	x	-	TT ^T _{GA} TACTAACa	-	y	-	T ^T _A T ^T aaT ^T _{AG}	↓
	(b)									A ^T _A ttc	

- (a) For references see Table II
- (b) a and t are used at positions with a preference for A and T




Fig. 5. Sequence analysis of the 5'-end of the L17a mRNA. cDNA synthesis was performed as described in Materials and Methods using the 73 nucleotides long primer depicted in Fig. 1. The arrows indicate the 5'-termini of the L17a mRNA; 'intron' indicates the splice junction. In b the concentration of ddNTP's was doubled.

signals found in higher eukaryotes (36,37), though in yeast the variability of the bases at the splice sites is even less. In addition to the sequences at the 5'- and 3'-termini of the introns another region of striking sequence homology occurs near the 3'-end. This region contains the TACTAAC-box which has been demonstrated to be essential for splicing in yeast (34,38,39). Apart from these primary structure elements no other homologous sequences nor common secondary structures can be observed in the introns of yeast ribosomal protein genes.

Table IV. Transcription initiation sites of several yeast ribosomal protein genes.

S10 (a)	ATCCAGTGTGAATAGACGACTGAGCCATC ATG
L16 (b)	GAACCAAGAACATACAACATAGCCAAAG ATG
S33 (c)	GAAAACCAAGCTAGCAATC ATG
L17a	CAAAACCAACACACCTATATATAACTAATA ATG
L25	TAAACAAAGAAGATCAATAAGATAAA ATG

(a) R.J. Leer, unpublished results
 (b) (24)
 (c) (12)




Fig. 7. Mapping of the 3'-ends of the L17a and L25 mRNAs. Lanes a-e show the product analysis using the 316 nucleotides long probe for the L17a gene (see Materials and Methods and Fig. 1). Lanes f-i show the product analysis using the 264 nucleotides long probe for the L25 gene (see Fig. 2). Lane a and b: 30 µg polyA containing yeast mRNA with 25 and 100 U S1 nuclease respectively; lanes c and d: 30 µg *B. licheniformis* RNA as a control with 25 and 100 U S1 nuclease respectively; lanes e and f: untreated probes; lanes g and h: 30 µg polyA-containing yeast mRNA with 25 and 100 U S1 nuclease respectively; lane i: 30 µg *B. licheniformis* RNA as a control with 25 U S1 nuclease. The GATC-ladder of a known sequence was used to measure the length of the S1-protected products.

the location of the intron boundaries in both ribosomal protein genes. The 5'-untranslated parts of the L17a and L25 mRNAs are AT-rich and contain no or only a few G-residues respectively. This feature of the leader region holds for most yeast ribosomal protein mRNAs. No obvious secondary structure elements such as may be involved in an autogenous control of translation in *E. coli* (41) could be detected. The transcription start sites of the L17a and L25 genes map in a region near the sequence PyAAPu, which has been proposed as a common capping site for RNA polymerase B transcripts (42). The data summarized in Table IV illustrate that AACCAA forms an additional conserved element in some but not all ribosomal protein genes. Finally the L17a and L25 mRNAs have an A at position -3, consistent with the preferred surroundings of the initiation codon observed in other eukaryotic mRNAs (43) and moreover use the 5'-proximal AUG as the start codon thus fitting into the ribosome-scanning model for translation initiation (43).

The 3'-ends of the L17a and L25 mRNAs were mapped by S1 nuclease analysis. The results presented in Fig. 7 show the mature transcript to terminate at 103 ± 10 nucleotides from the stop codon for the L17a mRNA and at approx. 80 nucleotides from the stopcodon for the L25 mRNA. The observed heterogeneity of the 3'-ends to our opinion reflects the actual situation; it may be the result of processing events, and is not due to the experimental conditions used in the mapping experiments. The trailer regions of the two genes contain a striking sequence homology: a tripartite common element ATTAT---T₄CT₄---ACAAT can be recognized. A similar sequence is present in the untranslated 3'-terminal region of the rp29 mRNA (35), but it does not occur in any other yeast ribosomal protein genes so far investigated. Whether this common sequence fulfils a function in transcription or translation remains to be examined. In this respect it is notable that the 3'-terminal regions of the L25 and L17a genes do not contain sequences that are identical to the consensus sites proposed to be involved in transcription termination in yeast (44). The sequence AATAAA, which may act as a polyadenylation signal (45), is present 57 nucleotides downstream from the stop codon of the L25 gene but it is not found in the trailer region of the L17a gene.

A comparative analysis of the 5'-flanking regions of several yeast ribosomal protein genes has revealed a number of more or less conserved sequence elements that, however, are not common to all ribosomal protein genes (46). These boxes may function in the coordinate control of ribosomal protein gene expression. We searched for similar sequences in the upstream regions of the genes encoding L17a and L25, (cf. Table V). The most striking homologous

Table V. Homologous sequences upstream of yeast ribosomal protein genes (a).

	consensus	L17a	L25
HOMOL1	AACATC ^{TG} _{CA} TCA (approx. -300)	AACATCGTACA (-351)	not found
HOMOL2	^T _A CATCTNTA (approx. -300)	ACATCTGTA (-350)	not found
HOMOL3	TC ^T _{GG} CTTC ^T _C T (approx. -150)	not found	GGCTTCCC (-143) GCCCTCCT (-225)
HOMOL4	^T _A AT ^T _A TINCA (varying positions)	TATTTTICA (-303)	TATTTIGCA (-440)
HOMOL5	TATT ^{TT} _{AA} (approx. -80)	TATTTT (-60)	TATTTT (-83)

(a) the consensus sequences for the various boxes are from Ref. 46.

sequences, HOMOL1 and HOMOL2, are present at -351 and -350 respectively in front of the L17a gene, but these boxes are absent from the gene encoding L25. On the other hand HOMOL3 is present at two positions, -143 and -225, upstream of the L25 gene but is not found in the L17a gene. HOMOL4 can be recognized in the 5'-flanking sequences of both ribosomal protein genes though at a rather long distance from the respective initiation codons. HOMOL5 which resembles the TATA box (47) and therefore may fulfil a role in initiation of transcription by RNA polymerase B, occurs at -60 and -83 from the ATG codons of the L17a and L25 genes respectively. Finally the upstream region of the L17a gene contains a CT-rich sequence, which has been suggested to define a high-efficiency yeast promoter (48). A CT-block occurs only occasionally in front of yeast ribosomal protein genes, in addition e.g. those encoding S33 (12) and L29 (32). The significance of all presumed promoter and regulatory sequences mentioned above has to be established by deletion experiments and site-directed mutagenesis. These studies are currently performed.

ACKNOWLEDGEMENTS

The authors are indebted to Dr. H.A. Raué for critical reading of the manuscript and to Mrs. P.G. Brink for preparing the typescript. This work was supported in part by the Netherlands Foundation for Chemical Research (S.O.N.) with financial aid from the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).

REFERENCES

1. Dean, D. and Nomura, M. (1982) in *The Cell Nucleus* (Busch, H. and Rothblum, L. eds) Vol XII, pp. 185-212, Acad. Press, New York.
2. Woolford, J.L., Hereford, L.H. and Rosbash, M. (1979) *Cell* 18, 1247-1259.
3. Fried, H.M., Pearson, N.J., Kim, C.H. and Warner, J.R. (1981) *J. Biol. Chem.* 256, 10176-10183.
4. Bollen, G.H.P.M., Cohen, L.H., Mager, W.H., Klaassen, A.W. and Planta, R.J. (1981) *Gene* 14, 279-182.
5. Fried, H.M. and Warner, J.R. (1982) *Nucl. Acids Res.* 10, 3133-3148.
6. Schultz, L.D. and Friesen, J.D. (1983) *J. Bact.* 155, 8-14.
7. Laskin, J.C. and Woolford, J.L. (1983) *Nucl. Acids Res.* 11, 403-420.
8. Bollen, G.H.P.M., Molenaar, C.M.T., Cohen, L.H., Van Raamsdonk-Duin, M.M.C., Mager, W.H. and Planta, R.J. (1982) *Gene* 18, 29-37.
9. Molenaar, C.M.T. (1984) Ph.D. Thesis, Free University, Amsterdam.
10. Woolford, J.L. and Rosbash, M. (1981) *Nucl. Acids Res.* 9, 5021-5028.
11. Rosbash, M., Harris, P.K.W., Woolford, J.L. and Teem, J.L. (1981) *Cell* 24, 679-686.
12. Leer, R.J., Van Raamsdonk-Duin, M.M.C., Schoppink, P.J., Cornelissen, M.T.E., Cohen, L.H., Mager, W.H. and Planta, R.J. (1983) *Nucl. Acids Res.* 11, 7759-7768.
13. Ng, R. and Abelson, J. (1980) *Proc. Natl. Acad. Sci. USA* 77, 3912-3916.
14. Gallwitz, D. and Sures, I. (1980) *Proc. Natl. Acad. Sci. USA* 77, 2546-2550.
15. Miller, A.M. (1984) *The EMBO J.* 3, 1061-1065.
16. Gorenstein, C. and Warner, J.R. (1976) *Proc. Natl. Acad. Sci. USA* 73, 1547-1551.
17. Warner, J.R. and Gorenstein, C. (1977) *Cell* 11, 201-212.
18. Kim, C.K. and Warner, J.R. (1983) *Mol. Cell. Biol.* 3, 457-465.
19. Pearson, N.J., Fried, H.M. and Warner, J.R. (1982) *Cell* 29, 347-355.
20. Smith, H.O. and Birnstiel, M.L. (1976) *Nucl. Acids Res.* 3, 2387-2398.
21. Messing, J. (1982) in *Genetic Engineering/Principles and Methods* (Setlow, J.K. and Hollaender, A. eds) Plenum Press, New York, London, Vol 4, pp. 19-34.
22. Sanger, F., Coulson, A.R., Barrell, B.G., Smith, A.J.H. and Roe, B.A. (1980) *J. Mol. Biol.* 143, 161-178.
23. Maxam, A.M., and Gilbert, W. (1977) *Proc. Natl. Acad. Sci. USA* 74, 560-564.
24. Leer, R.J., Van Raamsdonk-Duin, M.M.C., Mager, W.H. and Planta, R.J. (1984) *FEBS Lett.*, submitted.
25. Itoh, T., Higo, K.-J. and Otaka, E. (1979) *Biochemistry* 18, 5787-5793.
26. Bollen, G.H.P.M., Mager, W.H. and Planta, R.J. (1981) *Mol. Biol. Rep.* 8, 37-44.
27. El-Baradi, T.T.A.L., Raué, H.A., De Regt, V.C.H.F. and Planta, R.J. (1984) *Eur. J. Biochem.*, submitted.
28. Bennetzen, J.L. and Hall, B.D. (1982) *J. Biol. Chem.* 257, 3026-3031.
29. Leer, R.J., Van Raamsdonk-Duin, M.M.C., Molenaar, C.M.T., Cohen, L.H., Mager, W.H. and Planta, R.J. (1982) *Nucl. Acids Res.* 10, 5869-5878.
30. Teem, J.L. and Rosbash, M. (1983) *Proc. Natl. Acad. Sci. USA* 80, 4403-4407.
31. Schaap, P.J., Molenaar, C.M.T., Mager, W.H. and Planta, R.J. (1984) *Curr. Genet.*, submitted.
32. Käufer, N.F., Fried, H.M., Schwindinger, W.F., Jasin, M. and Warner, J.R. (1983) *Nucl. Acids Res.* 11, 3123-3135.
33. Ammerer, G., Hitzeman, R., Hagie, F., Barta, A. and Hall, B.D. (1981) in *Recombinant DNA* (Walton, A.G. ed) Elsevier Scientific Publishing Co., Amsterdam, pp. 185-197.
34. Pikielny, C.W., Teem, J.L. and Rosbash, M. (1983) *Cell* 34, 395-403.
35. Mitra, G. and Warner, J.R. (1984) *J. Biol. Chem.*, in press.
36. Breathnach, R. and Chambon, P. (1981) *Ann. Rev. Biochem.* 50, 349-383.
37. Mount, S.M. (1982) *Nucl. Acids Res.* 10, 459-527.

38. Langford,C.J. and Gallwitz,D. (1983) *Cell* 33, 519-527.
39. Langford,C.J., Klinz,F.J., Donath,C. and Gallwitz,D. (1984) *Cell* 36, 645-653.
40. Thill,G.P., Kramer,R.A., Turner,K.J. and Bostian,K.A. (1983) *Mol. Cell. Biol.* 3, 570-579.
41. Olins,P.O. and Nomura,M. (1981) *Nucl. Acids Res.* 9, 1757-1764.
42. Burke,R.L., Tekamp-Olson,P. and Najarian,R. (1983) *J. Biol. Chem.* 258, 2193-2201.
43. Kozak,M. (1981) *Nucl. Acids Res.* 9, 5233-5252.
44. Zaret,K.S. and Sherman,F. (1982) *Cell* 28, 563-573.
45. Fitzgerald,M. and Shenk,T. (1981) *Cell* 24, 251-260.
46. Teem,J.L., Abovich,N., Käufer,N.F., Schwindinger,W.F., Warner,J.R., Levy,A., Woolford,J., Leer,R.J., Van Raamsdonk-Duin,M.M.C., Mager,W.H., Planta,R.J., Schultz,L., Friesen, J.D. and Rosbash,M. (1984) *Nucl. Acids Res.*, submitted.
47. Benoist,C. and Chambon,P. (1981) *Nature* 290, 304-310.
48. Dobson,M.J., Tuite,M.E., Roberts,N.A., Kingsman,A.J., Kingsman,S.M., Perkins,R.E., Conroy,S.C., Dunbar,B and Fothergrill,L.A. (1982) *Nucl. Acids Res.* 10, 2625-2637.